

JAX® Mice, Clinical & Research Services (JMCRS)

Production Health Report

Area: MP13

HEALTH STATUS

| ledow | PATHOGEN & OPPORTUNIST FREE |
|------------|-----------------------------|
| \bigcirc | PATHOGEN FREE |

BARRIER LEVEL

| \odot | MAXIMUM BARRIER |
|------------|------------------|
| \bigcirc | HIGH BARRIER |
| \bigcirc | STANDARD BARRIER |

Test Results: #positive/#tested

Please consult our website for descriptions of our health statuses and barrier levels.

PATHOGENS AND OTHER ORGANISMS- EXCLUDED FROM ALL BARRIERS (SHIPPING STOPPED)

If one of these organisms is found in any JMCRS area, all shipments are suspended and customers are notified.*

| | | | | rest Results: #positive/#tested | | | | | |
|--|--------------------|-------------|-------------|---------------------------------|------------|------------|-----------|-----------------------|--|
| Organism | Sample Tested | Test Method | Frequency | Nov 4 '24 | Sep 23 '24 | Aug 12 '24 | Jul 1' 24 | Previous 12 months | |
| VIRUSES | | | | | | | | | |
| Ectromelia virus | Serum | MFI | 6 weeks | 0/16 | 0/19 | 0/19 | 0/18 | 0/146 | |
| GDVII (Theiler's) virus | Serum | MFI | 6 weeks | 0/16 | 0/19 | 0/19 | 0/18 | 0/146 | |
| Hantaan virus | Serum | MFI | 6 weeks | 0/16 | 0/19 | 0/19 | 0/18 | 0/146 | |
| K virus | Serum | ELISA | annually | - | - | - | - | 0/16 | |
| LDH elevating virus (LDEV) | Serum | PCR | annually | - | - | - | - | 0/10 | |
| Lymphocytic choriomeningitis (LCMV) | Serum | MFI | 6 weeks | 0/16 | 0/19 | 0/19 | 0/18 | 0/146 | |
| Mouse adenovirus (MAV) | Serum | MFI | 6 weeks | 0/16 | 0/19 | 0/19 | 0/18 | 0/146 | |
| Mouse cytomegalovirus (MCMV) | Serum | MFI | 6 weeks | 0/16 | 0/19 | 0/19 | 0/18 | 0/146 | |
| Murine chapparvovirus (MuCPV) | Cecum or feces | PCR | 6 weeks | 0/10 | 0/08 | 0/08 | 0/09 | 0/75 | |
| Mouse hepatitis virus (MHV) | Serum | MFI | 6 weeks | 0/16 | 0/19 | 0/19 | 0/18 | 0/146 | |
| Mouse minute virus (MMV) | Serum | MFI | 6 weeks | 0/16 | 0/19 | 0/19 | 0/18 | 0/145 | |
| Mouse norovirus (MNV) | Serum | MFI | 6 weeks | 0/16 | 0/19 | 0/19 | 0/18 | 0/146 | |
| Mouse parvovirus (MPV) | Serum | MFI | 6 weeks | 0/16 | 0/19 | 0/19 | 0/18 | 0/146 | |
| Mouse parvovirus (MPV)† | Lymph node | PCR | 6 weeks | 0/10 | 0/08 | 0/08 | 0/09 | 0/82 | |
| Mouse thymic virus (MTV) | Serum | IFA | quarterly | - | - | 0/19 | 0/18 | 0/64 | |
| Pneumonia virus of mice (PVM) | Serum | MFI | 6 weeks | 0/16 | 0/19 | 0/19 | 0/18 | 0/146 | |
| Polyoma virus | Serum | ELISA | annually | - | - | - | - | 0/16 | |
| Reovirus 3 (REO 3) | Serum | MFI | 6 weeks | 0/16 | 0/19 | 0/19 | 0/18 | 0/146 | |
| Rotavirus (EDIM) | Serum | MFI | 6 weeks | 0/16 | 0/19 | 0/19 | 0/18 | 0/145 | |
| Sendai virus | Serum | MFI | 6 weeks | 0/16 | 0/19 | 0/19 | 0/18 | 0/146 | |
| BACTERIA & MYCOPLASMA | | | | | | | | | |
| Bordetella spp. | Oropharynx | Culture | 6 weeks | 0/27 | 0/27 | 0/27 | 0/27 | 0/234 | |
| Citrobacter rodentium | Intestine or feces | Culture | 6 weeks | 0/87 | 0/89 | 0/92 | 0/169 | 0/1849 | |
| Clostridium piliforme | Serum | ELISA | quarterly | - | - | 0/19 | 0/18 | 0/64 | |
| Corynebacterium bovis †¶ | Skin | PCR | 6 weeks | 0/09 | 0/07 | 0/07 | 0/07 | 0/64 | |
| Corynebacterium kutscheri | Oropharynx | Culture | 6 weeks | 0/27 | 0/27 | 0/27 | 0/27 | 0/234 | |
| Filobacterium rodentium (CAR bacillus) | Serum | MFI | 6 weeks | 0/16 | 0/19 | 0/19 | 0/18 | 0/146 | |
| Mycoplasma pulmonis | Serum | MFI | 6 weeks | 0/16 | 0/19 | 0/19 | 0/18 | 0/146 | |
| Mycoplasma spp. | Lung | PCR | semi-annual | - | - | 0/06 | - | 0/12 | |
| Salmonella spp. | Intestine or feces | Culture | 6 weeks | 0/87 | 0/89 | 0/92 | 0/169 | 0/1849 | |
| Streptobacillus moniliformis | Oropharynx | Culture | 6 weeks | 0/27 | 0/27 | 0/27 | 0/27 | 0/234 | |

^{*}Additional details regarding our health monitoring program and shipping policy.

 $^{{}^{\}dagger}{}$ The indicated tests are only performed in rooms that house immunodeficient mice.

Test Results: #positive/#tested

| Organism | Sample Tested | Test Method | Frequency | Nov 4 '24 | Sep 23 '24 | Aug 12 '24 | Jul 1' 24 | Previous 12 months |
|--|--------------------|-------------|-------------|-----------|------------|------------|-----------|--------------------|
| PARASITES | | | | | | | | |
| Encephalitozoon cuniculi | Serum | MFI | 6 weeks | 0/16 | 0/19 | 0/19 | 0/18 | 0/146 |
| Ectoparasites (fleas, lice, mites) | Fur | Visual | 6 weeks | 0/06 | 0/06 | 0/06 | 0/06 | 0/59 |
| Endoparasites (tapeworms, | Intestine or cecum | Visual | 6 weeks | 0/06 | 0/06 | 0/06 | 0/06 | 0/59 |
| pinworms§, and other helminths) | | | | | | | | |
| Follicle mites | Subcutis | Visual | 6 weeks | 0/27 | 0/27 | 0/27 | 0/27 | 0/234 |
| Protozoa (e.g., Giardia, Spironucleus, | Intestine | Microscopy | 6 weeks | 0/06 | 0/06 | 0/06 | 0/06 | 0/59 |
| etc.) | mtestine | | | | | | | |
| Toxoplasma gondii ‡ | Serum | ELISA | semi-annual | - | - | 0/06 | - | 0/12 |

OPPORTUNISTIC ORGANISMS MONITORED (SHIPPING NOT STOPPED)

All of these organisms are excluded from JMCRS maximum and high barriers, and most are excluded from standard barrier areas. When a confirmed finding of an excluded organism is made, an investigation is undertaken to identify and eliminate all infected mice from the barrier. Positive results- including results from investigations- are noted in this report, but shipping from the area is not suspended.*

| Organism | Sample Tested | Test Method | Frequency | Nov 4 '24 | Sep 23 '24 | Aug 12 '24 | Jul 1' 24 | Previous 12 months |
|-----------------------------------|---------------------|-------------|-------------|-----------|------------|------------|-----------|-----------------------|
| Dermatophytes | Skin swabs | Culture | semi-annual | - | - | 0/06 | - | 0/12 |
| Helicobacter spp. | Intestine or feces | PCR | 6 weeks | 0/10 | 0/08 | 0/08 | 0/09 | 0/75 |
| Klebsiella pneumoniae | Oropharynx, | Culture | 6 weeks | 0/87 | 0/89 | 0/92 | 0/169 | 0/1849 |
| | intestine, or feces | | | | | -, | | |
| Klebsiella oxytoca | Oropharynx, | Culture | 6 weeks | 0/87 | 0/89 | 0/92 | 0/169 | 0/1849 |
| | intestine, or feces | | | | | | | |
| Pasteurella multocida | Oropharynx | Culture | 6 weeks | 0/27 | 0/27 | 0/27 | 0/27 | 0/234 |
| Rodentibacter pneumotropicus / | Oropharynx | Culture | 6 weeks | 0/27 | 0/27 | 0/27 | 0/27 | 0/234 |
| Rodentibacter heylii | Oropilarytix | Culture | 0 WEEKS | | | | | |
| Pneumocystis murina † | Lung | PCR | 6 weeks | 0/10 | 0/08 | 0/08 | 0/09 | 0/82 |
| Proteus mirabilis | Oropharynx, | Culture | 6 weeks | 0/87 | 0/89 | 0/92 | 0/169 | 0/1849 |
| Froteus mirubinis | intestine, or feces | | | | | | | |
| Pseudomonas aeruginosa | Oropharynx, | Culture | 6 weeks | 0/87 | 0/89 | 0/92 | 0/169 | 0/1849 |
| - seudomonas deraginosa | intestine, or feces | | | | | | | |
| Staphylococcus aureus | Oropharynx | Culture | 6 weeks | 0/27 | 0/27 | 0/27 | 0/27 | 0/234 |
| Streptococcus pneumoniae | Oropharynx | Culture | 6 weeks | 0/27 | 0/27 | 0/27 | 0/27 | 0/234 |
| Beta-hemolytic Streptococcus spp. | 0 1 | Culture | 6 weeks | 0/27 | 0/27 | 0/27 | 0/27 | 0/234 |
| (non-group D) | Oropharynx | | | | | | | |
| Trichomonads | Intestine | Microscopy | 6 weeks | 0/06 | 0/06 | 0/06 | 0/06 | 0/59 |
| Yersinia enterocolitica | Oropharynx, | Culture | 6 weeks | 0/87 | 0/89 | 0/92 | 0/169 | 0/1849 |
| | intestine, or feces | | | | | | | |
| Yersinia pseudotuberculosis | Oropharynx, | Culture | 6 weeks | 0/87 | 0/89 | 0/92 | 0/169 | 0/1849 |
| | intestine, or feces | Culture | | | | | | |

[†]The indicated tests are only performed in rooms that house immunodeficient mice.

All tests were performed by The Jackson Laboratory

MP13

The Jackson Laboratory 600 Main Street Bar Harbor, ME 04609 1-800-422-6423

Rosalinda A. Doty DVM, PhD, DACVPDirector, Diagnostic and Pathology Services

[¶] One sample represents testing of 60 pooled animals

[§] Testing for this organisms is also performed by PCR

[‡]This testing is performed by an outside vendor